

Contingency & Crisis Strategies for Mask Usage

Based on CDC guidelines effective 4/2/2020

If your facility is facing a shortage of masks, the following CDC guidelines should be utilized to conserve masks (Contingency Strategy) while maintaining your protection:

- Implement limited re-use of facemasks: <u>use the same facemask by one provider</u> for multiple encounters with different patients
 - o Remove only if leaving a patient care area
- Do not touch outer surfaces of the mask during care, if you accidently touch your mask, immediately perform hand hygiene
 - Facemasks should be carefully folded inward; the outer surface is held against itself to reduce contact with the outer surface during storage.
 - o The folded mask is be <u>stored between uses in a clean sealable paper bag</u> or breathable container
 - o Label your bag or container so another provider does not accidently use
- The facemask should be removed and discarded if soiled, damaged, or hard to breathe through.
- Leave patient care area if you need to remove the facemask.

When no facemasks are available (Crisis Strategy), you have two options to protect yourself:

- Wear a face shield that covers the entire front
 - o It should extend to the chin and sides of the face with no facemask.
- Wear homemade masks, bandana or scarf for care of patients with COVID-19 as a last resort.
 - Homemade masks alone are not considered PPE, their capability to protect is unknown.
 - Homemade masks can be used as a cover for extended use surgical masks to extend their use. The homemade mask can be laundered daily
 - Homemade masks should ideally be used in combination with a face shield that covers the entire front (that extends to the chin or below) and sides of the face.

Other PPE tips

- Reserve PPE for direct patient care & use PPE per CDC guidance
- To prevent self-contamination during PPE use, please don and doff correctly, folding inward when removing and then performing hand hygiene after properly disposing

Remember: all patients with known or suspected COVID-19, with ACTIVE symptoms, will be placed on a medical hold until their symptoms have resolved and their physician has cleared them to resume therapy services.